

## TABLES

**Table 1.** Baseline characteristics according to clinical and virological outcome of 1061 patients treated with HCQ+AZ  $\geq 3$  days at IHU Méditerranée infection Marseille, France with day 0 between March 3 and March 31, 2020.

	Poor virological outcome <sup>a</sup>	Good outcome	Poor clinical outcome <sup>a,b</sup>	Total
	n (%)	n (%)	n (%)	n (%)
<b>Group size</b>	47 (4.4%)	973 (91.7%)	46 (4.3%)	1061 (100%)
<b>Age (years)</b>				
Mean (SD)	47.9 (17.5)	42.4 (14.7)	69.2 (14.0)	43.6 (15.6)
Median [Min-Max]	<b>48.0 [18.0-89.0]*</b>	42.0 [14.0-86.0]	<b>69.0 [31.0-95.0]***</b>	43.0 [14.0-95.0]
<b>Male</b>	19 (40.4%)	450 (46.3%)	23 (50%)	492 (46.4)
<b>Chronic condition(s) and treatment(s)</b>				
<b>Chronic conditions</b>				
Cancer	0 (0.0%)	21 (2.2%)	<b>7 (15.2%)***</b>	28 (2.6%)
Diabetes	3 (6.4%)	66 (6.8%)	<b>9 (19.6%)***</b>	78 (7.4%)
Coronary artery disease	2 (4.3%)	36 (3.7%)	<b>9 (19.6%)***</b>	46 (4.3%)
Hypertension	8 (17%)	120 (12.3%)	<b>23 (50.0%)***</b>	149 (14%)
Chronic respiratory diseases	8 (17%)	96 (9.9%)	8 (17.4%)	111 (10.5%)
Obesity	1 (2.1%)	57 (5.9%)	4 (8.7%)	62 (5.8%)
<b>Comedication(s)</b>				
Biguanides (metformin)	1 (2.1%)	15 (1.5%)	<b>4 (8.7%)**</b>	20 (1.9%)
Selective beta blocking agents	<b>6 (12.8%)**</b>	22 (2.3%)	<b>9 (19.6%)***</b>	34 (3.2%)
Dihydropyridine derivatives	3 (6.4%)	23 (2.4%)	<b>8 (17.4%)***</b>	34 (3.2%)
Angiotensin II receptor blockers	<b>6 (12.8%)**</b>	22 (2.3%)	<b>14 (30.4%)***</b>	40 (3.8%)
HMG CoA reductase inhibitors	4 (8.5%)	28 (2.9%)	<b>7 (15.2%)***</b>	38 (3.6%)
Diuretics	2 (4.3%)	28(2.9%)	<b>5 (10.9%)*</b>	35(3.3%)
<b>Time between onset of symptoms and first day of treatment start (days)<sup>e</sup></b>				
Mean (SD)	4.3 (2.5)	6.5 (3.9)	5.9 (4.0)	6.4 (3.8)
Median [Min-Max]	<b>4.0 [0.0-9.0]***</b>	6.0 [0.0-27.0]	<b>5.0 [0.0-16.0]***</b>	6.0 [0.0-27.0]
<b>Clinical classification (NEWS score)</b>				
0 – 4 (low)	<b>43 (91.5%)*</b>	948 (97.4%)	<b>19 (41.3%)***</b>	1008 (95.0%)
5 – 6 (medium)	2 (4.3%)	14 (1.4%)	10 (21.7%)	25 (2.4%)
$\geq 7$ (high)	2 (4.3%)	11 (1.1%)	17 (37.0%)	28 (2.6%)
<b>Low-dose pulmonary CT-scanner within 72 hours of admission<sup>d</sup></b>				
Normal	11/37 (29.7%)	231/642 (36.0%)	<b>4/39 (10.3%)***</b>	245/714 (34.3%)
Limited	23/37 (62.2%)	277/642 (43.2%)	10/39 (25.6%)	307/714 (43.0%)
Medium	3/37 (8.1%)	123/642 (19.2%)	20/39 (51.3%)	146/714 (20.5%)
Severe	0/37 (0.0%)	11/642 (1.7%)	5/39 (12.8%)	16/714 (2.2%)
<b>Viral load at inclusion (Ct - nasal)<sup>e</sup></b>				
Mean (SD)	23.4 (5.1)	26.8 (4.9)	25.6 (4.8)	26.6 (5.0)
Median [Min-Max]	<b>22.1 [14.8-34.0]***</b>	27.3 [12.8-34.0]	25.8 [15.0-33.2]	27.0 [12.8-34.0]
<b>Hydroxychloroquine levels at day 2 (<math>\mu\text{g/mL}</math>)<sup>f</sup></b>				
Mean (SD)	0.25 (0.17)	0.26 (0.16)	0.20 (0.17)	0.25 (0.16)
Median [Min-Max]	0.19 [0.07-0.70]	0.22 [0.00-1.01]	<b>0.15 [0.00-0.75]**</b>	0.21 [0.00-1.01]
Number $\leq 0.1\mu\text{g/mL}$	4/24 (16.7%)	15/206 (7.3%)	<b>12/37 (32.4%)***</b>	30/263 (11.4%)

Poor virological outcome (PVirO): viral shedding persistence at day 10; Poor clinical outcome (PClinO): either death or transfer to intensive care unit (ICU) or hospitalization for 10 days or more; Good outcome: individuals who belonged neither to the PClinO group nor the PVirO group. SD: standard deviation. <sup>a</sup>Five patients belonged to both the PVirO and PClinO outcome so the sum of frequencies may be above 1061. <sup>b</sup>Including 8 deaths. <sup>c</sup>Data available for 928 patients (56 patients who did not declare any symptom before treatment start were excluded and 77 with missing data), <sup>d</sup>for 714 patients, <sup>e</sup>for 992 patients and <sup>f</sup>for 263 patients. On low-dose pulmonary CT-scanner, patients were classified as no involvement (lack of lung involvement (ground glass opacities, consolidation or crazy paving pattern); minimal involvement (subtle ground glass opacities); intermediate involvement (less than 50% of segment involvement in no more than 5 segments) and severe involvement (involvement of more than 5 segments). The denominator was mentioned when the result was not available for all patients. \*  $p < 0.05$ ; \*\*  $p < 0.01$ ; \*\*\*  $p < 0.001$  (Fisher's exact test, Student t-test, Wilcoxon-Mann-Whitney where appropriate; reference group is good outcome).